

12

**EUROPEAN PATENT APPLICATION**

21 Application number: 79105421.6

22 Date of filing: 31.12.79

51 Int. Cl.<sup>3</sup>: **C 07 D 521/00, A 61 K 31/00**  
 // C07D249/12, C07D249/14,  
 C07D207/22, C07D233/84,  
 C07D235/28, C07D239/56,  
 C07D207/36, C07D277/16,  
 C07D239/38, C07D237/18,  
 C07D239/36, C07D277/74,  
 C07D401/04, C07D235/06,  
 C07D285/12

30 Priority: 12.01.79 CH 309/79  
 08.08.79 CH 7274/79

43 Date of publication of application: 06.08.80  
 Bulletin 80/16


84 Designated Contracting States: AT BE CH DE FR GB IT  
 LU NL SE

71 Applicant: **SANDOZ AG, Lichtstrasse 35, CH-4002 Basel (CH)**

72 Inventor: **Berner, Heinz, Dr., Martinstrasse 83, A-1180 Wien (AT)**  
 Inventor: **Turnowsky, Friederike, Dr., Wienerbruckstrasse 87, A-2344 Maria Enzersdorf (AT)**  
 Inventor: **Laber, Georg, Dr., Alserstrasse 18/30, A-1090 Wien (AT)**  
 Inventor: **Hildebrandt, Johannes, Dr., St. Laurentgasse 12, A-2512 Oeynhausen (AT)**

54 New pleuromutilin derivatives, their production and use.

57 Novel pleuromutilin derivatives of formula I,



In which

R<sub>1</sub> is ethyl or vinyl,

m is 0 or 1, and

R<sub>2</sub> is a heterocyclic radical, in which a 5- or 6-membered, unsaturated or saturated heterocyclic ring containing one or more hetero atoms selected from oxygen, sulphur and nitrogen, is attached to the -S(CH<sub>2</sub>)<sub>m</sub>-group, provided that when n is 0, R<sub>2</sub> is other than pyridyl,

their production and use as antimicrobial agents are described.


**EP 0 013 768 A1**

0013768

Case 900-9240

NEW PLEUROMUTILIN DERIVATIVES, THEIR PRODUCTION AND USE

This invention provides compounds of formula I,



in which  $R_1$  is ethyl or vinyl,

$m$  is 0 or 1, and

$R_2$  is a heterocyclic radical, in which a 5-


5

or 6-membered, unsaturated or saturated heterocyclic ring containing one or more hetero atoms selected from oxygen, sulphur and nitrogen, is attached to the  $-S(CH_2)_m-$  group,

10

provided that when  $m$  is 0,  $R_2$  is other than pyridyl.

The invention also provides a process for the production of compounds of formula I, characterised by reacting a compound of formula II,




II

in which  $R_1$  is as defined above, and

5  $R_5$  is chlorine, bromine or  $-\text{OSO}_2R_7$ , in which  
 $R_7$  is alkyl or aryl,

with a compound of formula III,



III

in which  $m$  and  $R_2$  are as defined above.




The process is suitably carried out in the presence  
 10 of an alkali metal lower alkoxide, for example sodium  
 ethoxide or methoxide. This is preferably produced in  
situ. Conveniently, the compound of formula III may be  
 dissolved in a solution of sodium in a water-free lower  
 alkanol, e.g. methanol or ethanol. A solution of the  
 15 compound of formula II in an inert organic solvent, e.g.  
 an aliphatic ketone, such as methyl ethyl ketone or acet-  
 one, is then conveniently added. The process is suitably  
 effected at a temperature from room temperature to the

reflux temperature of the reaction mixture, in particular from 22° to 55°C. The reaction time may typically vary from 2 to 12 hours.

The resulting compounds of formula I may be isolated and purified using conventional techniques. Where required, free base forms thereof may be converted into salt forms, in particular into acid addition salt and quaternary ammonium salt forms, in conventional manner, and vice versa.

10  $R_2$  suitably signifies a 5- or 6-membered saturated or unsaturated heterocyclic ring containing one or more hetero atoms selected from oxygen, sulphur and nitrogen. The ring may be unsubstituted. Alternatively, it may be mono- or poly-substituted. Suitable substituents include  
15 mercapto, thioxo, hydroxy, lower alkyl, lower alkanoyl, lower sulfoxyl, nitro, lower alkylsulphonyl, trifluoromethyl, formyl, lower alkoxycarbonyl, lower hydroxyalkyl, lower dihydroxyalkyl and halogen.

Further suitable substituents are of formulae II,  
20 III and IV,



in which either  $R_3$  and  $R_4$  are the same or different  
and each is hydrogen, lower hydroxyalkyl,  
lower dihydroxyalkyl, unsubstituted or  
substituted lower alkanoyl, lower alkyl  
5 sulfonyl or lower alkyl,  
or  $R_3$  and  $R_4$  together with the nitrogen atom  
form a piperaziny radical, which may be  
substituted on the second nitrogen atom by  
lower alkyl, lower hydroxyalkyl or lower  
10 dihydroxyalkyl,  
 $n$  is 2 to 5,  
 $X$  is oxygen or sulphur,  
and  $R_6$  is lower alkyl or lower alkocycarbonyl.

Other substituents include further 5- or 6-membered,  
15 saturated or unsaturated heterocyclic rings, e.g. pyridyl,  
which may themselves be unsubstituted or mono- or poly-  
substituted as described above.

Finally, the heterocyclic ring of  $R_2$  may suitably  
be fused to one or more 5- or 6-membered, saturated or  
20 unsaturated carbocyclic or heterocyclic, preferably carbo-  
cyclic, e.g. benzene, rings. This ring may also be unsub-  
stituted or similarly mono- or poly-substituted, as des-  
cribed above.

As used herein, the term "lower" signifies preferably  
25 of 1 to 4, more preferably 1 to 2 carbon atoms. "Halogen"  
signifies chlorine, bromine, fluorine or iodine, prefer-

ably chlorine, bromine or fluorine, more preferably chlorine.


The preferred heterocyclic rings in  $R_2$  linked to the  $-S(CH_2)_m-$  radical contain one or more hetero atoms selected from nitrogen and sulphur. The more preferred ring contains at least one nitrogen atom.

One group of such hetero rings may contain nitrogen as the sole hetero atom, in particular 1, 2 or 3 nitrogen hetero atoms. Suitable 5- or 6-membered hetero rings containing a single nitrogen atom include pyridine (when  $m$  is 1), pyrrole and 4,5-dihydro-3H-pyrrole. Suitable 5- or 6-membered rings containing 2 nitrogen atoms include imidazole, pyridazine, pyrimidine. Such rings may be fused to, e.g. one or more benzene rings, e.g. to form benzimidazole or perimidine. Suitable 5- or 6-membered hetero rings containing 3 nitrogen atoms include 1,2,4-triazole.

Another group of hetero rings may contain 1 nitrogen atom and 1 sulphur atom, e.g. thiazole, 4,5-dihydrothiazole and benzothiazole. Another group of hetero rings contain 2 nitrogen and 1 sulphur atom, e.g. 1,3,4-thiadiazole.

Preferred compounds are those in which the heterocyclic ring of  $R_2$  is bound to the  $-S-(CH_2)_m-$  group via a ring carbon atom. Particularly preferred compounds are those in which the hetero ring of  $R_2$  is 1,2,4-triazole.

A particularly preferred group of compounds are those in which  $R_2$  is formula V,



V

in which X is hydrogen, lower alkylsulphonyl, amino,  
a group of formula II, III or IV; formyl,  
5 and Y is hydrogen, amino, trifluoromethyl, lower  
alkyl, a group of formula II, or pyridyl.

The compounds of formula I are indicated for use as  
chemotherapeutic agents, in particular as antimicrobial  
agents, as indicated, e.g. by their inhibiting effect against  
10 various bacterial strains, e.g. Staph. aureus, Staph.  
epidermis, Strept. pyogenes, Strept. aranson, Strept.  
pneumoniae, Strept. faecelis, Strept. viridans, Corynebact.  
pyogenes, Sarcina lutea, Klebsiella pneumoniae, and  
Haemophilus influenzae, in vitro in the series dilution  
15 test at a concentration, for example, of 0.01 to 25  $\mu\text{g/ml}$ ,  
and in in vivo tests in mice. The compounds also show an  
inhibiting effect against various mycoplasma, e.g. M. hom-  
inis, M. arthritidis, M. pneumoniae, and urea plasma  
urealyticum, and chlamydia, in vitro in the series dilu-  
20 tion test at concentrations of, for example, 0.008 to  
2.5  $\mu\text{g/ml}$ .

The compounds also show an inhibiting effect against various obligatory anaerobes, e.g. *Bacteroides fragilis*, *Bacteroides melaninogenicus*, *Sphaerophorus necrophorus*, *Clostridium perfringens*, etc., in vitro in the series  
5 dilution test at concentrations of for example 0.1 to 4 ug/ml, and in vivo in mice at a dosage of for example 50 to 200 mg/kg of animal body weight, p.o or s.c.

The compounds are therefore indicated for use as anti-microbial agents, in particular as antibacterially active  
10 antibiotics and for treatment of infections caused by obligatory anaerobes.

For the above-mentioned uses, an indicated suitable daily dosage is from about 1 to 3 g, suitably administered in divided dosages of from two to four times daily, or in  
15 retard form.

The compounds alone or in admixture with a tetracyclin may be administered orally or parenterally in such forms as tablets, capsules, powders, granulates, and injectable or



infusion preparations, e.g. solutions or suspensions.

The compounds may also be employed in the form of creams or tinctures. For veterinary purposes, the compounds may also be administered as food or drink additives.

5        It has also been found that mixtures of the compounds of formula I with a tetracyclin with R-factor coded tetracyclin resistance show a synergistic antibacterial effect against resistant strains of this type. This indicated for example by determination of the minimum inhibition concentration of the mixture and the  
10        individual components in the series dilution test, and by evaluating the results by the method of Löwe (isobole diagram), Die Antibiotika, Volume 1, part 1, 65ff, 1962. Conventional tetracyclines, e.g. chlorotetracyclin, oxy-  
15        tetracyclin, demethyltetracyclin, tetracyclin dioxycyclin, monocylin, metacyclin, and rolitetracyclin, may be employed in such mixtures. The quantity of the compound of formula I in such mixtures is suitably 10 to 90%, preferably 20 to 35%, in particular 25%, while the quantity of  
20        the tetracyclin is suitably from 90 to 10%, preferably 80 to 65%, particularly 75% (these percentages being by weight).

      The mixtures are particularly indicated in treating infections of the gastrointestinal tract and other local  
25        infections of the organ.

The compounds of formula I, when used alone or in admixture with a tetracyclin, may be employed in free base form or in the form of chemotherapeutically acceptable acid addition or quaternary ammonium salts. These  
5 salt forms have the same order of activity as the free base forms.

Suitable acid addition salt forms include the hydrochloride, hydrogen fumarate, fumarate and naphthalene-1,5-sulphonate.

10 The compounds (or mixtures thereof with a tetracycline) may be admixed with a chemotherapeutically acceptable diluent or carrier and, optionally other conventional excipients for the production of galenic forms. Suitable excipients include sweeteners, aromas, colouring  
15 agents, preserving agents, e.g. ethyl-o-hydroxybenzoate, fillers or carriers, e.g. diluents, such as calcium carbonate, disintegrating agents, e.g. starch or alginic acid, binding agents, e.g. starch, gelatine or acacia, and lubricating agents, e.g. magnesium stearate, stearic acid or  
20 talc. Oral liquid forms may contain conventional suspending agents, e.g. methylcellulose, tragacanth or sodium alginate. Suitable wetting agents include lecithin, polyoxyethane stearate and polyoxyethylene sorbitan monooleate. For the production of capsules, suitable diluents include  
25 calcium carbonate, calcium phosphate and kaolin.

0013768

- 10 -

900-9240

The preferred compound of formula I is that of Example 1, hereinafter.

The following Examples illustrate the invention. All temperatures are in degrees Centigrade.

EXAMPLE 1: 19,20-Dihydro-14-O-[(3-amino-1,2,4-triazol-  
5-yl)thioacetyl]mutilin

2.3 g of sodium are taken up in 500 ml of absolute ethanol. After formation of the sodium ethoxide, 11.6 g of 3-amino-5-mercapto-1,2,4-triazol are added to the solution.

The mixture is allowed to react for 3 hours at 25° and is then mixed with a solution of 53.5 g of 19,20-dihydro-22-O-tosyl-pleuromutilin in 200 ml of ethylmethyl-  
10 ketone. The homogenous reaction mixture is held for 12 hours at 25° and then poured onto water and extracted 3 times with 500 ml of ethyl acetate. The purified ethyl acetate extract is shaken with water, dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated in vacuo. The crude product is chromato-  
15 graphed over silica gel (eluant: ethyl acetate) to obtain the heading compound, m.p. 213-215° (isopropanol/H<sub>2</sub>O).

NMR (DMSO): 5.76 (broad, 2H, NH<sub>2</sub>); 5.52 (d, 1H, H<sub>14</sub>,

$J_{H_{14}H_{13}} = 8 \text{ Hz}$ );

3.76 (s, 2H, S-CH<sub>2</sub>-CO); 3.35 (m, 1H, H<sub>11</sub>).

20 IR (-KBr): 2600-3600 (broad), 1720, 1635, 1280 cm<sup>-1</sup>.

The compounds of the following Examples may be produced in manner analogous to that of Example 1, using appropriate starting materials in approximately equivalent amounts.

Example 2: 14-O-[3 Amino-1,2,4-triazol-5-yl]thioacetyl]-  
mutilin

---

NMR ( $\text{CDCl}_3$ ): 5.68 (d, 1H,  $\text{H}_{14}$ ,  $J_{\text{H}_{14}\text{H}_{13}} = 8 \text{ Hz}$ ); 3.34 (d, 1H,  $\text{H}_{11}$ ,  $J_{\text{H}_{11}\text{H}_{10}} = 6 \text{ Hz}$ ); 3.7 (s, 2H,  $\text{S-CH}_2\text{-CO}$ );

5 5.26-4.95 (m, 4H,  $\text{NH}_2 + 2\text{H}_{20}$ ); 6.5-6.16 (m, 1H,  $\text{H}_{19}$ ).

IR (KBr): 3300 (broad), 1720, 1625, 1575, 1270  $\text{cm}^{-1}$ .

EXAMPLE 3: 14-O-[(Imidazol-2-yl)thioacetyl]mutilin

---

10 NMR ( $\text{CDCl}_3/\text{DMSO } 5:1$ ): 6.98 (s, 2H, Imidazol H); 5.65 (d, 1H,  $\text{H}_{14}$ ,  $J_{\text{H}_{14}\text{H}_{13}} = 8 \text{ Hz}$ ); 3.76 (s, 2H,  $\text{S-CH}_2\text{CO}$ ); 3.4-3.2 (m, 1H,  $\text{H}_{11}$ ).

IR (KBr): 3600-2600 (broad), 1730, 1705, 1270  $\text{cm}^{-1}$ .

EXAMPLE 4: 14-O[(Perimidin-2-yl)thioacetyl]mutilin

---

15 NMR ( $\text{CDCl}_3$ ): 7.0-7.2 (m, 4H, arom. H); 5.74 (d, 1H,  $\text{H}_{14}$ ,  $J_{\text{H}_{14}\text{H}_{13}} = 8 \text{ Hz}$ ); 3.72 (AB-System, 2H,  $\text{S-CH}_2\text{-CO}$ ,  $J=16.2 \text{ Hz}$ ); 3.42-3.22 (m, 1H,  $\text{H}_{11}$ ).

IR (KBr): 3600-3100 (broad), 1720, 1625, 1585, 1270, 820, 770  $\text{cm}^{-1}$ .

EXAMPLE 5: 14-O-[(4,5-Dihydro-3H-pyrrol-2-yl)thio-  
acetyl]mutilin

---

NMR (CDCl<sub>3</sub>): 5.74 (d, 1H, H<sub>14</sub>, J<sub>H<sub>14</sub>H<sub>13</sub></sub> = 8 Hz),  
3.96-3.58 (m, 4H, 2-Pyrrolidin H + S-CH<sub>2</sub>-CO),  
5 3.36 (dd, 1H, H<sub>11</sub>, J=6.3 Hz, J=10.8 Hz).  
IR (CHCl<sub>3</sub>): 1720, 1590 cm<sup>-1</sup>.

EXAMPLE 6: 14-O-[(Benzimidazol-2-yl)thioacetyl]mutilin

---

NMR (CDCl<sub>3</sub>): 7.5-7.0 (m, 4H, arom. H); 5.64 (d, 1H, H<sub>14</sub>,  
J<sub>H<sub>14</sub>H<sub>13</sub></sub> = 8.2 Hz); 4.1 (s, 2H, S-CH<sub>2</sub>-CO);  
10 3.38 (d, 1H, H<sub>11</sub>, J<sub>H<sub>11</sub>H<sub>10</sub></sub> = 6.3 Hz).  
IR (KBr): 3550-2600 (breit), 1720, 1270, 740 cm<sup>-1</sup>.

EXAMPLE 7: 14-O-[(4-Methylsulfonyl-5-amino-1,2,4-  
triazol-3-yl)thioacetyl]mutilin

---

NMR (CDCl<sub>3</sub>): 5.9 (s, 2H, NH<sub>2</sub>); 5.76 (d, 1H, H<sub>14</sub>,  
15 J<sub>H<sub>14</sub>H<sub>13</sub></sub> = 8 Hz); 3.81 (s, 2H, S-CH<sub>2</sub>-CO);  
3.3 (s, 3H, CH<sub>3</sub>SO<sub>2</sub>-); 3.4 (m, 1H H<sub>11</sub>).  
IR (KBr): 3400 (broad), 1720, 1625, 1265, 1275,  
1180 cm<sup>-1</sup>.

EXAMPLE 8: 14-O-[(3-Mercaptopyridazin-6-yl)thioacetyl]-  
20 19,20-dihydromutilin

---

NMR (CDCl<sub>3</sub>): 7.5 (d, 1H, arom. H, J=9 Hz); 6.9 (d, 1H,  
arom. H, J=9 Hz); 5.64 (d, 1H, H<sub>14</sub>, J<sub>H<sub>14</sub>H<sub>13</sub></sub> =  
8 Hz); 3.79 (s, 2H, S-CH<sub>2</sub>-CO); 3.44 (d,  
1H, H<sub>11</sub>, J<sub>H<sub>11</sub>H<sub>10</sub></sub> = 6 Hz).

IR (KBr): 3400 (broad), 1720, 1270, 1155, 1140  $\text{cm}^{-1}$ .

EXAMPLE 9: 14-O-[(2-Isopropyl-4-hydroxypyrimidin-6-yl)methylthioacetyl]mutilin

---

5 NMR ( $\text{CDCl}_3$ ): 6.32 (s, 1H, arom. H); 5.81 (d, 1H,  $\text{H}_{14}$ ,  $J_{\text{H}_{14}\text{H}_{13}} = 8 \text{ Hz}$ ); 3.62 (s, 2H,  $\text{S-CH}_2\text{-CO}$ ); 3.3 (s, 2H,  $\text{S-CH}_2\text{-Arom.}$ ); 3.4 (m, 1H,  $\text{H}_{11}$ ).

IR (KBr): 3400 (broad), 1720, 1650, 1590, 1275  $\text{cm}^{-1}$ .

10 EXAMPLE 10: 14-O-[[3-(4-Hydroxyäthylpiperazin-1-yl)äthylthio]pyridazin-6-yl]thioacetyl]mutilin hydrochloride form

---

NMR ( $\text{CDCl}_3$ /  
 $\text{CD}_3\text{OD}$  50:1): 7.26 (s, 2H, arom. H); 5.7 (d, 1H,  $\text{H}_{14}$ ,  $J_{\text{H}_{14}\text{H}_{13}} = 8 \text{ Hz}$ ).

IR (KBr): 3350 (broad), 1720, 1380, 1270, 1140  $\text{cm}^{-1}$ .

15 EXAMPLE 11: 14-O-[(6-Nitrobenzothiazol-2-yl)thioacetyl]mutilin

---

20 NMR ( $\text{CDCl}_3$ ): 8.68 (d, 1H, arom. H,  $J=2.3 \text{ Hz}$ ); 8.3 (dd, 1H, arom. H,  $J_1=2.3 \text{ Hz}$ ,  $J_2=9 \text{ Hz}$ ); 7.84 (d, 1H, arom. H,  $J=9 \text{ Hz}$ ); 5.78 (d, 1H,  $\text{H}_{14}$ ,  $J_{\text{H}_{14}\text{H}_{13}} = 8 \text{ Hz}$ ); AB-System: ( $\nu_A=3.18$ ,  $\nu_B=3.06$ ,  $\text{S-CH}_2\text{-CO}$ ,  $J=16.2 \text{ Hz}$ ); 3.34 (dd, 1H,  $\text{H}_{11}$ ,  $J=6 \text{ Hz}$ ,  $J=10.8 \text{ Hz}$ ).

IR (KBr): 3400 (broad), 1720, 1510, 1325, 1265, 1010  $\text{cm}^{-1}$ .

EXAMPLE 12: 14-O-[(4-Methyl-6-hydroxypyrimidin-2-yl)-  
thioacetyl]mutilin

---

NMR (DMSO): 5.94 (s, 1H, arom. H); 5.53 (d, 1H, H<sub>14</sub>,  
J<sub>H<sub>14</sub>H<sub>13</sub></sub> = 8 Hz); 3.9 (s, 2H, S-CH<sub>2</sub>CO);  
5 3.4 (m, 1H, H<sub>11</sub>); 2.14 (s, 3H, CH<sub>3</sub>).

IR (KBr): 3400 (broad), 1720, 1650, 1525, 1270, 1160 cm<sup>-1</sup>.

EXAMPLE 13: 14-O-[(4-Ethoxycarbonyl-3,5-dimethylpyrrol-2-  
yl)thioacetyl]mutilin

---

NMR (CDCl<sub>3</sub>): 8.75 (b, 1H, NH); 5.74 (d, 1H, H<sub>14</sub>, J<sub>H<sub>14</sub>H<sub>13</sub></sub> =  
10 8 Hz); 4.26 (q, 2H, -OCH<sub>2</sub>CH<sub>3</sub>); 3.22 (s, 2H,  
S-CH<sub>2</sub>CO); 3.36 (dd, 1H, H<sub>11</sub>, J=6.3 Hz, J=10.8 Hz);  
2.46 (s, 3H, CH<sub>3</sub>-Pyrrol); 2.28 (s, 3H, CH<sub>3</sub>-  
Pyrrol).

IR (KBr): 3600-3200 (broad), 1720, 1780, 1250, 1100 cm<sup>-1</sup>.

15 EXAMPLE 14: 14-O-[(1-Methylimidazol-2-yl)thioacetyl]-  
mutilin

---

NMR (CDCl<sub>3</sub>): 7.0 (d, 1H, ImidazoleH, J=1.8 Hz); 6.86  
(d, 1H, ImidazoleH, J=1.8 Hz); 5.7 (d, 1H,  
H<sub>14</sub>, J<sub>H<sub>14</sub>H<sub>13</sub></sub> = 8 Hz); 3.78 (s, 2H, S-CH<sub>2</sub>CO);  
20 3.62 (s, 3H, N-CH<sub>3</sub>); 3.34 (m, 1H, H<sub>11</sub>).

IR (KBr): 3200 (broad), 1720, 1270 cm<sup>-1</sup>.

m.p.: 135-136°.



EXAMPLE 15: 14-O-[(3-Mercaptopyridazin-6-yl)thioacetyl]-  
mutilin

---

NMR (CDCl<sub>3</sub>): AB-System of the Pyridazine protons ( $\nu_A=7.5$ ,  
 $\nu_B=6.9$ ,  $J_{AB}=9$  Hz); 5.64 (1H, H<sub>14</sub>,  $J_{H_{14}H_{13}}=7$  Hz);  
5 3.78 (s, 2H, S-CH<sub>2</sub>-CO); 3.44 (1H, H<sub>11</sub>,  $J_{H_{11}H_{10}}=6.3$  Hz).

IR (KBr): 3400 broad (OH), 1725 (CO), 1140, 1155 cm<sup>-1</sup>.

EXAMPLE 16: 14-O-[(3-Chlorpyridazin-6-yl)thioacetyl]-  
mutilin

---

10 NMR (CDCl<sub>3</sub>): 5.78 (d, 1H, H<sub>14</sub>,  $J_{H_{14}H_{13}}=8$  Hz); 3.38  
(m, 1H, H<sub>11</sub>); 1.44 (s, 3H, (CH<sub>3</sub>)<sub>15</sub>);  
1.21 (s, 3H, (CH<sub>3</sub>)<sub>18</sub>);  
AB-System of the Pyridazin-H ( $\nu_A=7.29$ ,  $\nu_B=7.33$ ,  
 $J_{AB}=9$  Hz);  
15 AB-System (CH<sub>2</sub>)<sub>22</sub> ( $\nu_A=4.12$ ,  $\nu_B=4.02$ ,  $J_{AB}=16$  Hz).

IR (KBr): 3500 (OH), 1720 (CO) cm<sup>-1</sup>.

EXAMPLE 17: 14-O-[(4,5-Dihydrothiazol-2-yl)thioacetyl]-  
mutilin

---

20 NMR (CDCl<sub>3</sub>): 5.75 (d, 1H, H<sub>14</sub>,  $J_{H_{14}H_{13}}=8$  Hz); 3.81  
(s, 2H, -(CH<sub>2</sub>)<sub>22</sub>-); 1.46 (s, 3H, (CH<sub>3</sub>)<sub>15</sub>);  
1.17 (s, 3H, (CH<sub>3</sub>)<sub>18</sub>); 4.14 (t, 2H, -S-CH<sub>2</sub>,  
 $J=8$  Hz); 3.4 (t, 2H, =N-CH<sub>2</sub>).

IR (KBr): 3500 (OH), 1710 (CO), 1570 cm<sup>-1</sup>.

EXAMPLE 18: 14-O-[(3-Diethylaminoäthylthiopyridazin-6-yl)thioacetyl]mutilin hydrogen fumarate form

---

NMR (CDCl<sub>3</sub>): 7.14 (s, 2H, Pyridazin-H); 5.78 (d, 1H, H<sub>14</sub>,  $J_{H_{14}H_{13}} = 8$  Hz); 4.1 (s, 2H, S-CH<sub>2</sub>-CO); 3.4 (m, 3H, H<sub>11</sub> und CH<sub>2</sub>-N<); 2.9 (m, 2H, CH<sub>2</sub>-S-); 2.66 (q, 4H, N-CH<sub>2</sub>-CH<sub>3</sub>); 1.1 (t, 6H, N-CH<sub>2</sub>-CH<sub>3</sub>); 1.46 (s, 3H, (CH<sub>3</sub>)<sub>15</sub>); 1.1 (s, 3H (CH<sub>3</sub>)<sub>18</sub>).

IR (KBr): 3400 (broad, OH), 1720 (CO), 1140 cm<sup>-1</sup>.

10 EXAMPLE 19: 14-O-[(4-Amino-1,2,4-triazol-3-yl)-thioacetyl]mutilin

---

NMR (CDCl<sub>3</sub>): 8.26 (s, 1H, Triazol-H); 5.7 (d, 1H, H<sub>14</sub>,  $J_{H_{14}H_{13}} = 8$  Hz); 5.08 (s, 2H, NH<sub>2</sub>), AB-System ( $v_A = 3.86$ ,  $v_B = 3.75$ ,  $J_{AB} = 16.2$  Hz, S-CH<sub>2</sub>-O); 3.34 (dd, 1H, H<sub>11</sub>,  $J = 6.3$  Hz,  $J = 10.2$  Hz).

IR (KBr): 3400 (broad), 1720 cm<sup>-1</sup>.

EXAMPLE 20: 14-O-[(3-(4-Pyridyl)-1,2,4-triazol-5-yl)-thioacetyl]mutilinhydrochlorid

---

20 NMR (DMSO): 8.95 (d, 2H, Pyridin-H,  $J = 6.3$  Hz); 8.36 (d, 2H-Pyridin-H,  $J = 6.3$  Hz); 5.55 (d, 1H, H<sub>14</sub>,  $J_{H_{14}H_{13}} = 8$  Hz); 4.16 (s, 2H, S-CH<sub>2</sub>-CO); 3.4 (d, 1H, H<sub>11</sub>,  $J_{H_{10}H_{11}} = 6.3$  Hz).

IR (KBr): 3600-2500 (broad), 1725, 1635 cm<sup>-1</sup>.

EXAMPLE 21: 14-O-[(4-Amino-3-trifluoromethyl-1,2,4-  
 triazol-5-yl)thioacetyl]mutilin

NMR (CDCl<sub>3</sub>): 5.74 (d, 1H, H<sub>14</sub>, J<sub>H<sub>14</sub>H<sub>13</sub></sub> = 8 Hz); 5.18  
 (s, 2H, NH<sub>2</sub>); 3.9 (s, 2H, S-CH<sub>2</sub>CO); 3.38  
 (m, 1H, H<sub>11</sub>).

IR (KBr): 3400 (broad), 1720, 1190, 1150 cm<sup>-1</sup>.

EXAMPLE 22: 14-O-[(4-Amino-3-methyl-1,2,4-triazol-  
 5-yl)thioacetyl]mutilin

NMR (CDCl<sub>3</sub>): 5.72 (d, 1H, H<sub>14</sub>, J<sub>H<sub>14</sub>H<sub>13</sub></sub> = 8 Hz); 4.97  
 (s, 2H, NH<sub>2</sub>); AB-System (v<sub>A</sub>=3.84, v<sub>B</sub>=3.69,  
 J<sub>AB</sub>=16.2 Hz, S-CH<sub>2</sub>-CO); 3.38 (dd, H<sub>11</sub>, J=6.3 Hz,  
 J=10.2 Hz).

IR (KBr): 3400 (broad), 1725 cm<sup>-1</sup>.

EXAMPLE 23: 14-O-[(3-Methyl-4-acetamido-1,2,4-triazol-  
 5-yl)thioacetyl]mutilin

NMR (CDCl<sub>3</sub>): 5.7 (d, 1H, H<sub>14</sub>, J<sub>H<sub>14</sub>H<sub>13</sub></sub> = 8 Hz); 3.8 (s,  
 2H, S-CH<sub>2</sub>-CO); 3.38 (m, 1H, H<sub>11</sub>); 2.33  
 (s, 3H, CH<sub>3</sub>CO-N); 2.26 (s, 3H, Triazol-CH<sub>3</sub>).

IR (KBr): 3400 (broad), 1720, 750 cm<sup>-1</sup>.

EXAMPLE 24: 14-O[(3-(Methoxysulfonylethylcarboxamido)-  
 1,2,4-triazol-5-yl)thioacetyl]mutilin

NMR (CDCl<sub>3</sub>): 5.72 (d, 1H, H<sub>14</sub>, J<sub>H<sub>14</sub>H<sub>13</sub></sub> = 8 Hz); 3.82

(s, 2H, S-CH<sub>2</sub>-CO); 3.34 (m, 1H, H<sub>11</sub>);  
3.14 (s, 3H, -O-CH<sub>3</sub>).

IR (KBr): 3400 (broad), 1720, 1625, 1550, 1305,  
1110, 730 cm<sup>-1</sup>.

5 EXAMPLE 25: 14-O-[(1-Ethylaminocarbonyl-3-amino-1,2,4-  
triazol-5-yl)thioacetyl]mutilin

NMR (CDCl<sub>3</sub> /

• CD<sub>3</sub>OD 10:1): 5.74 (d, 1H, H<sub>14</sub>, J<sub>H<sub>14</sub>H<sub>13</sub></sub> = 8 Hz); 3.75  
(s, 2H, S-CH<sub>2</sub>-CO); 3.38 (q, 2H, CH<sub>3</sub>-CH<sub>2</sub>-N);  
3.4 (m, 1H, H<sub>11</sub>); 1.26 (t, 3H, CH<sub>3</sub>-CH<sub>2</sub>-N).

10 IR (KBr): 3540, 3430, 3310, 1710, 1630, 1300 cm<sup>-1</sup>.

m.p.: 230-232°.

EXAMPLE 26: 14-O-[(3-Amino-4-formyl-1,2,4-triazol-  
5-yl)thioacetyl]mutilin

15 NMR (CDCl<sub>3</sub>): 8.52 (s, 1H, Formyl-H); 5.74 (d, 1H, H<sub>14</sub>,  
J<sub>H<sub>14</sub>H<sub>13</sub></sub> = 8 Hz); 3.82 (s, 2H, S-CH<sub>2</sub>CO);  
3.36 (m, 1H, H<sub>11</sub>).

IR (KBr): 3600-2800 (broad), 1725, 1585, 1290 cm<sup>-1</sup>.

EXAMPLE 27: 14-O-[(3-Amino-1-(carboethoxythiocarbamyl)-  
1,2,4-triazol-5-yl)thioacetyl]mutilin

20 NMR (CDCl<sub>3</sub>): 7.74 (b, 2H, NH<sub>2</sub>); 5.81 (d, 1H, H<sub>14</sub>, J<sub>H<sub>14</sub>H<sub>13</sub></sub> = 8 Hz);

4.37 (q, 2H, O-CH<sub>2</sub>CH<sub>3</sub>); 3.83 (s, 2H, S-CH<sub>2</sub>-CO); 3.4 (d, 1H, H<sub>11</sub>, J<sub>H<sub>11</sub>H<sub>10</sub></sub> = 6.3 Hz); 1.38 (t, 3H, O-CH<sub>2</sub>CH<sub>3</sub>).

IR (KBr): 3300 (broad), 1770, 1725, 1635, 1465, 1185 cm<sup>-1</sup>.

5

EXAMPLE 28: 14-O-[(3 Amino-4-(Ethylaminothiocarbonyl)-1,2,4-triazol-5-yl)thioacetyl]mutilin

---

NMR (CDCl<sub>3</sub>): 8.56 (m, 1H, NH); 7.4 (b, 2H, NH<sub>2</sub>); 5.76 (d, 1H, H<sub>14</sub>, J<sub>H<sub>14</sub>H<sub>13</sub></sub> = 8 Hz); 3.78 (s, 2H, S-CH<sub>2</sub>-CO); 3.68 (m, 2H, N-CH<sub>2</sub>-CH<sub>3</sub>); 3.38 (dd, 1H, H<sub>11</sub>, J=6.3 Hz, J=10.2 Hz); 1.33 (t, 3H, N-CH<sub>2</sub>-CH<sub>3</sub>).

10

IR (KBr): 3340 (broad), 1720, 1630, 1290 cm<sup>-1</sup>.

EXAMPLE 29: 14-O-[[4-Bis-(methylsulfonylamino)-1,2,4-triazol-3-yl]thioacetyl]mutilin

---

15

NMR (CDCl<sub>3</sub>): 8.28 (s, 1H, Triazol-H); 5.72 (d, 1H, H<sub>14</sub>, J<sub>H<sub>14</sub>H<sub>13</sub></sub> = 8 Hz); AB-System (v<sub>A</sub>=4.18, v<sub>B</sub>=4.02, J<sub>AB</sub> = 16.2 Hz, S-CH<sub>2</sub>-CO); 3.6 (s, 3H, CH<sub>3</sub>SO<sub>2</sub>-); 3.58 (s, 3H, CH<sub>3</sub>SO<sub>2</sub>-); 3.34 (m, 1H, H<sub>11</sub>).

20

IR (KBr): 3450 (broad), 1720, 1380, 1160 cm<sup>-1</sup>.

EXAMPLE 30: 14-O-[(Benzimidazol-2-yl-methyl)thioacetyl]mutilinhydrochloride

---

NMR (CDCl<sub>3</sub>): 7.6 (m, 2H, arom.H); 7.2 (m, 2H, arom.H);

5.8 (d, 1H,  $H_{14}$ ,  $J_{H_{14}H_{13}} = 8$  Hz); 4.07  
 (s, 2H, S-CH<sub>2</sub>-Arom.); 3.42 (m, 1H,  $H_{11}$ );  
 3.25 (s, 2H, S-CH<sub>2</sub>-CO).

IR (KBr): 3600-2700 (broad), 1720, 1270, 740 cm<sup>-1</sup>.

5 EXAMPLE 31: 14-O-[(2-Methyl-4-hydroxypyrimidin-6-yl)-  
methylthioacetyl]mutilin

10 NMR (CDCl<sub>3</sub>): 6.3 (s, 1H, NH); 5.74 (d, 1H,  $H_{14}$ ,  $J_{H_{14}H_{13}} =$   
 8 Hz); 3.58 (s, 2H, S-CH<sub>2</sub>-Arom.); 3.36  
 (m, 1H,  $H_{11}$ ); 3.14 (s, 2H, S-CH<sub>2</sub>-CO); 2.46  
 (s, 3H, Pyrimidin-CH<sub>3</sub>).

IR (KBr): 3400 (broad), 1720, 1650, 1590, 1270, 1110 cm<sup>-1</sup>.

EXAMPLE 32: 19,20-Dihydro-14-O-[(3-diethylaminoethyl-  
thiopyridazin-6-yl)thioacetyl]mutilin, fumarate  
form

15 NMR (CDCl<sub>3</sub>): 7.14 (s, 2H, Pyridazin-H); 6.78 (s, 1H,  
 Fumarsäure-H); 5.6 (d, 1H,  $H_{14}$ ,  $J_{H_{14}H_{13}} =$   
 8 Hz); AB-System ( $v_A=4.14$ ,  $v_B=3.92$ ,  $J_{AB}=$   
 16.2 Hz); 2.97 (q, 4H, N-CH<sub>2</sub>-CH<sub>3</sub>);  
 1.23 (t, 6H, N-CH<sub>2</sub>-CH<sub>3</sub>); 3.6-3.1 (m, 4H,  
 S-CH<sub>2</sub>-CH<sub>2</sub>-N<).

20 IR (KBr): 3400 (breit), 1720, 1385, 1140, 1110 cm<sup>-1</sup>.

EXAMPLE 33: 14-O-{[3-(2-Pyridyl)-1,2,4-triazol-5-yl]-  
thioacetyl}mutilin

NMR (CDCl<sub>3</sub>): 8.82 (d, 1H, Pyridin-H, J=5 Hz);

8.24 (d, 1H, Pyridin-H,  $J=10$  Hz), 7.9 (m, 1H, Pyridin-H), 7.45 (m, 1H, Pyridin-H), 5.78 (d, 1H,  $H_{14}$ ,  $J_{H_{14}H_{13}} = 8$  Hz), 4.0 (s, 2H, S-CH<sub>2</sub>-CO), 3.4 (m, 1H,  $H_{11}$ ).

5 IR (KBr): 3500-2800 (broad), 1720, 1450, 1280 cm<sup>-1</sup>.

UV (CH<sub>3</sub>OH): 232 nm ( $\epsilon = 12400$ ), 252 (8170).

EXAMPLE 34: 14-O-[(2-Methyl-1,3,4-thiadiazol-5-yl)thioacetyl]-  
mutilin

10 NMR (CDCl<sub>3</sub>): 5.78 (d, 1H,  $H_{14}$ ,  $J_{H_{14}H_{13}} = 8.1$  Hz), 4.07 (s, 2H, CH<sub>2</sub>-S-CO), 3.38 (dd, 1H,  $H_{11}$ ,  $J_{H_{11}H_{10}} = 6.3$  Hz,  $J_{H_{11}OH} = 10.8$  Hz), 2.72 (s, 3H, CH<sub>3</sub>-thiadiazol).

IR (KBr): 3400 (OH) (broad) 1730 (CO) cm<sup>-1</sup>.

UV (CH<sub>3</sub>OH): 264 nm ( $\epsilon=5330$ ).

15 EXAMPLE 35: 14-O[(2-Amino-1,3,4-thiadiazol-5-yl)thioacetyl]-  
mutilin


NMR (CDCl<sub>3</sub>): 5.74 (d, 1H,  $H_{14}$ ,  $J_{H_{14}H_{13}} = 8.1$  Hz), 5.32 (b, 2H, NH<sub>2</sub>), 3.88 (s, 2H, S-CH<sub>2</sub>-CO), 3.34 (m, 1H,  $H_{11}$ ).

IR (KBr): 3400 (NH<sub>2</sub>, OH), 1730 (CO) cm<sup>-1</sup>.

UV (CH<sub>3</sub>OH): 282 nm ( $\epsilon=7150$ ).

WHAT WE CLAIM IS:

1. A compound of formula I,



in which  $R_1$  is ethyl or vinyl,

$m$  is 0 or 1, and

$R_2$  is a heterocyclic radical, in which a 5-  
 or 6-membered, unsaturated or saturated  
 heterocyclic ring containing one or more  
 hetero atoms selected from oxygen, sulphur  
 and nitrogen, is attached to the  $-S(CH_2)_m-$   
 group,

provided that when  $m$  is 0,  $R_2$  is other  
 than pyridyl,




and acid addition and quaternary ammonium salts thereof.

2. The compound of Claim 1, which is 19,20-dihydro-14-O-[(3-amino-1,2,4-triazol-5-yl)thioacetyl]mutilin.

3. A chemotherapeutic composition comprising a compound of Claim 1, in association with a chemotherapeutically acceptable diluent or carrier.

4. A process for the production of a compound of formula I, stated in Claim 1, or an acid addition or quaternary ammonium salt thereof, which comprises reacting  
10 a compound of formula II,




II

in which  $R_1$  is as defined above, and

$R_5$  is chlorine, bromine or  $-\text{OSO}_2R_7$ , in which

$R_7$  is alkyl or aryl,

with a compound of formula III,



III

15 in which  $m$  and  $R_2$  are as defined above.

5. The steps, features, compositions and compounds referred to or indicated in the specification and/or claims of this application, individually or collectively, and any and all combinations or any two or more of said steps or features.



## EUROPEAN SEARCH REPORT

EP 79 10 5421

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl. *)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
	<p>GB - A - 1 312 148 (BIOCHEMIE)</p> <p>* Claims *</p> <p>--</p> <p>THE JOURNAL OF ANTIBIOTICS, vol. XXIX, no. 9, published September 1976, pages 915-922 Tokyo, JP. H. EGGER et al.: "New pleuromutillin derivatives with enhanced antimicrobial activity. I. Synthesis"</p> <p>* Page 917, formula 1-58; table I, compounds 23-35, 58; pages 919-920, Method D *</p> <p>--</p> <p>THE JOURNAL OF ANTIBIOTICS, vol. XXIX, no. 9, published September 1976, pages 923-927 Tokyo, JP. H. EGGER et al.: "New pleuromutillin derivatives with enhanced antimicrobial activity. II. Structure-activity correlations"</p> <p>* Page 923, formula 1-58; page 920, table I, compounds 23-35, 58; page 926 *</p> <p>----</p>	<p>1,3,5</p> <p>1,4,5</p> <p>1,3,5</p>	<p>C 07 D 521/00</p> <p>A 61 K 31/00//</p> <p>C 07 D 249/12</p> <p>249/14</p> <p>207/22</p> <p>233/84</p> <p>235/28</p> <p>239/56</p> <p>207/36</p> <p>277/16</p> <p>239/38</p> <p>237/18</p> <p>239/36</p> <p>277/74</p> <p>401/04-7</p> <p>TECHNICAL FIELDS SEARCHED (Int. Cl. *)</p> <p>CATEGORY OF CITED DOCUMENTS</p> <p>X: particularly relevant</p> <p>A: technological background</p> <p>O: non-written disclosure</p> <p>P: intermediate document</p> <p>T: theory or principle underlying the invention</p> <p>E: conflicting application</p> <p>D: document cited in the application</p> <p>L: citation for other reasons</p> <p>&amp;: member of the same patent family, corresponding document</p>
X	The present search report has been drawn up for all claims		
Place of search	Date of completion of the search	Examiner	
The Hague	31-03-1980	NUYTS	

0013768



European Patent  
Office

**EUROPEAN SEARCH REPORT**

Application number

EP 79 10 5421

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl. <sup>3</sup> )
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
			C 07 D 235/06 285/12
			TECHNICAL FIELDS SEARCHED (Int. Cl. <sup>3</sup> )